

## **Amendments to the Claims**

This listing of claims will replace prior versions and listings of claims in the application:

### **Listing of Claims**

1. (currently amended) A method of reducing post-surgical vomiting in a patient comprising administering to ~~[[a]]~~ the patient ~~undergoing general anesthesia~~ a therapeutically effective amount of Doxylamine Succinate and Pyridoxine Hydrochloride before general anesthesia is administered to the patient and before the patient presents symptoms of post-surgical vomiting.
2. (previously presented) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered before surgery.
3. (currently amended) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are also administered during surgery.
4. (currently amended) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are also administered after surgery.
5. (original) The method of claim 4, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered after surgery at regular intervals.
6. (original) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered before, during and after surgery.
7. (original) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered orally.
8. (original) The method of claim 7, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered orally in a delayed release formulation.
9. (original) The method of claim 8, wherein the orally delayed release formulation is enterically coated.

10. (original) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are formulated in a pharmaceutically acceptable carrier.
11. (original) The method of claim 1, further defined as administering at least about 10 mg of Doxylamine Succinate and at least about 10 mg of Pyridoxine Hydrochloride.
12. (original) The method of claim 1, further defined as administering at least about 20 mg of Doxylamine Succinate and at least about 20 mg of Pyridoxine Hydrochloride.
13. (canceled)
14. (original) The method of claim 1, wherein the patient is a woman.
15. (original) The method of claim 1, wherein the surgery is performed on an outpatient basis.
16. (currently amended) The method of claim 1, wherein Doxylamine Succinate and Pyridoxine Hydrochloride are administered on an evening prior to surgery, a morning of the day of surgery ~~or~~ and after surgery.
17. (original) The method of claim 1, wherein Doxylamine Succinate is administered before, at substantially the same time or after Pyridoxine Hydrochloride is administered to the patient.

Claims 18-24 (canceled)